

COUPLED PAINLEVÉ II SYSTEMS IN DIMENSION 4 AND THE SYSTEMS OF TYPE $A_4^{(1)}$

YUSUKE SASANO

ABSTRACT. We find and study coupled Painlevé II systems in dimension 4, which can be obtained by a degeneration from the systems of type $A_4^{(1)}$. We compare these systems with other types of coupled Painlevé II systems from the viewpoint of the local index. We also give the phase spaces for these systems.

0. INTRODUCTION

In 2001, K. Takano described the confluence process in certain phase spaces of Painlevé systems as deformations of manifolds [8]. In 1998, Noumi and Yamada [1] proposed the systems of type $A_4^{(1)}$, which can be considered as a 4-parameter family of fourth-order coupled Painlevé IV systems in dimension 4, whose Hamiltonians are given as follows:

$$\begin{aligned} H &= xy(2y - x - 2t) - 2\beta_1 y - \beta_2 x + zw(2w - z - 2t) - 2\beta_3 w - \beta_4 z + 4yzw \\ &= H_{IV}(x, y, t; \beta_1, \beta_2) + H_{IV}(z, w, t; \beta_3, \beta_4) + 4yzw. \end{aligned}$$

Here, x, y, z and w denote unknown complex variables, and $\beta_1, \beta_2, \beta_3$ and β_4 are complex parameters.

It is well-known that P_{IV} has a confluence to the second Painlevé equation P_{II} , where two accessible singularities come together into a single singularity. This suggests the possibility that there exists a procedure for searching for higher order versions of Painlevé II, by using Takano's description of the confluence process from P_{IV} to P_{II} for the coordinate systems (x, y) and (z, w) , respectively. In this vein, the goal of this work is to find a fourth-order version of the Painlevé II equation. We present a 2-parameter family of fourth-order algebraic ordinary differential equations which can be considered as coupled Painlevé II systems in dimension 4, and which is given as follows:

$$(1) \quad \left\{ \begin{array}{l} \frac{dx}{dt} = -x^2 + y + w - \frac{t}{2} \\ \frac{dy}{dt} = 2xy + \alpha_1 \\ \frac{dz}{dt} = -z^2 + y + w - \frac{t}{2} \\ \frac{dw}{dt} = 2zw + \alpha_2. \end{array} \right.$$

Here x, y, z and w denote unknown complex variables, and α_1 and α_2 are complex parameters.

Our differential system is equivalent to a Hamiltonian system. The Hamiltonian is given as follows:

$$\begin{aligned} H &= \frac{y^2}{2} - (x^2 + \frac{t}{2})y - \alpha_1 x + \frac{w^2}{2} - (z^2 + \frac{t}{2})w - \alpha_2 z + yw \\ &= H_{II}(x, y, t; \alpha_1) + H_{II}(z, w, t; \alpha_2) + yw. \end{aligned}$$

From the viewpoint of symmetry, it is interesting to point out that our differential system is invariant under the following transformations:

$$\begin{aligned} s_1 &: (x, y, z, w, t; \alpha_1, \alpha_2) \rightarrow (x + \frac{\alpha_1}{y}, y, z, w, t; -\alpha_1, \alpha_2), \\ s_2 &: (x, y, z, w, t; \alpha_1, \alpha_2) \rightarrow (x, y, z + \frac{\alpha_2}{w}, w, t; \alpha_1, -\alpha_2), \\ s_3 &: (x, y, z, w, t; \alpha_1, \alpha_2) \rightarrow \\ &\quad (x, \frac{-1+\alpha_1+\alpha_2+xy-yz}{x-z}, z, \frac{1-\alpha_1-\alpha_2+xw-zw}{x-z}, t; 1-\alpha_2, 1-\alpha_1), \\ \pi &: (x, y, z, w, t; \alpha_1, \alpha_2) \rightarrow (z, w, x, y, t; \alpha_2, \alpha_1). \end{aligned}$$

We regard the system (1) as an algebraic vector field v on $(x, y, z, w, t) \in \mathbb{C}^4 \times B$:

$$v = \frac{\partial}{\partial t} + \frac{dx}{dt} \frac{\partial}{\partial x} + \frac{dy}{dt} \frac{\partial}{\partial y} + \frac{dz}{dt} \frac{\partial}{\partial z} + \frac{dw}{dt} \frac{\partial}{\partial w}$$

with $B = \mathbb{C}$. If we take a relative compactification $\mathbb{P}^4 \times B$ of $\mathbb{C}^4 \times B$, the extended vector field $\tilde{v}$ satisfies the following conditions:

$$\tilde{v} \in H^0(\mathbb{P}^4, \Theta_{\mathbb{P}^4}(-\log \mathcal{H})(\mathcal{H})).$$

Here $\mathcal{H}$ is the boundary divisor in $\mathbb{P}^4$ and $\Theta_{\mathbb{P}^4}(-\log \mathcal{H})(\mathcal{H})$ is the subsheaf of $\Theta_{\mathbb{P}^4}$ whose local section v satisfies $v(f) \in (f)$ for any local equation f of $\mathcal{H}$. Let us extend the regular vector field v on $\mathbb{C}^4 \times B$ to a rational vector field $\tilde{v}$ on $\mathbb{P}^4 \times B$. Then $\tilde{v}$ has poles along the boundary divisor $\mathcal{H}$. Moreover, $\tilde{v}$ has accessible singularities along subvarieties in the boundary divisor $\mathcal{H}$. (For the definition of accessible singularities, see Definition 1.1.) In order to explain our main results, we recall the definition of a symplectic transformation and its properties (cf. [4]). Let

$$\begin{aligned} \varphi &: x = x(X, Y, Z, W, t), \quad y = y(X, Y, Z, W, t), \quad z = z(X, Y, Z, W, t), \\ &\quad w = w(X, Y, Z, W, t), \quad t = t \end{aligned}$$

be a biholomorphic mapping from a domain D in $\mathbb{C}^5 \ni (X, Y, Z, W, t)$ into $\mathbb{C}^5 \ni (x, y, z, w, t)$. We say that the mapping is symplectic if

$$dx \wedge dy + dz \wedge dw = dX \wedge dY + dZ \wedge dW,$$

where t is considered as a constant or a parameter, namely, if, for $t = t_0$, $\varphi|_{t=t_0} = \varphi|_{t=t_0}$ is a symplectic mapping from the t_0 -section D_{t_0} of D to $\varphi(D_{t_0})$. Suppose that the mapping is symplectic. Then any Hamiltonian system $dx/dt = \partial H/\partial y, dy/dt = -\partial H/\partial x, dz/dt = \partial H/\partial w, dw/dt = -\partial H/\partial z$ is transformed to $dX/dt = \partial K/\partial Y, dY/dt = -\partial K/\partial X, dZ/dt = \partial K/\partial W, dW/dt = -\partial K/\partial Z$, where

$$dx \wedge dy + dz \wedge dw - dH \wedge dt = dX \wedge dY + dZ \wedge dW - dK \wedge dt.$$

Here t is considered as a variable. By this equation, the function K is determined from H uniquely modulo functions of t , namely, modulo functions independent of X, Y, Z and W . For the vector field v in (1), we can prove the following theorem.

Theorem 0.1. *The phase space $\mathcal{X}$ over $B = \mathbb{C}$ for the vector field v in (1) is obtained by gluing nine copies of $\mathbb{C}^4 \times \mathbb{C}$:*

$$\begin{aligned} U_0 \times \mathbb{C} &= \mathbb{C}^4 \times \mathbb{C} \ni (x, y, z, w, t), \\ U_j \times \mathbb{C} &= \mathbb{C}^4 \times \mathbb{C} \ni (x_j, y_j, z_j, w_j, t) \quad (j = 1, 2, \dots, 8), \end{aligned}$$

via the following symplectic transformations :

- 1) $x_1 = \frac{1}{x}$, $y_1 = -x(xy + \alpha_1)$, $z_1 = z$, $w_1 = w$,
- 2) $x_2 = x$, $y_2 = y$, $z_2 = \frac{1}{z}$, $w_2 = -z(zw + \alpha_2)$,
- 3) $x_3 = x(1 + (y + w - 2x^2 - t)x - 2w(z - x) - \alpha_1 - \alpha_2)$, $y_3 = \frac{1}{x}$,
 $z_3 = x^2(z - x)$, $w_3 = \frac{w}{x^2}$,
- 4) $x_4 = z^2(x - z)$, $y_4 = \frac{y}{z^2}$, $z_4 = z(1 + (y + w - 2z^2 - t)z - 2y(x - z) - \alpha_1 - \alpha_2)$, $w_4 = \frac{1}{z}$,
- 5) $x_5 = \frac{1}{x}$, $y_5 = -x(xy + \alpha_1)$, $z_5 = \frac{1}{z}$, $w_5 = -z(zw + \alpha_2)$,
- 6) $x_6 = x(1 + (y + w - 2x^2 - t)x + 2(xw - zw - \alpha_2) - \alpha_1 + \alpha_2)$, $y_6 = \frac{1}{x}$,
 $z_6 = \frac{81}{x^2(x - z)}$, $w_6 = \frac{x^2(x - z)(xw - zw - \alpha_2)}{81}$,
- 7) $x_7 = \frac{81}{z^2(z - x)}$, $y_7 = \frac{z^2(z - x)(yz - xy - \alpha_1)}{81}$,
 $z_7 = z(1 + (y + w - 2z^2 - t)z + 2(yz - xy - \alpha_1) + \alpha_1 - \alpha_2)$, $w_7 = \frac{1}{z}$,
- 8) $x_8 = -((x - z)y + \alpha_1 + \alpha_2 - 1)y$, $y_8 = \frac{1}{y}$, $z_8 = z$, $w_8 = w + y$.

Because every coordinate transformation is symplectic, the Hamiltonian system H in $U_0 \times \mathbb{C}$ is also written as a Hamiltonian system in each $U_j \times \mathbb{C}$ ($j = 1, 2, \dots, 7$). By direct calculation, we can verify the following theorem.

Theorem 0.2. *On the affine open set $(x_i, y_i, z_i, w_i, t) \in U_i \times B$ in Theorem 0.1, each Hamiltonian H_i on $U_i \times B$ is expressed as a polynomial in x_i, y_i, z_i, w_i and t and satisfies the following conditions :*

$$dx \wedge dy + dz \wedge dw - dH \wedge dt = dx_i \wedge dy_i + dz_i \wedge dw_i - dH_i \wedge dt.$$

This paper is organized as follows. In Section 1, we consider the relation between the systems (1) and the systems of type $A_4^{(1)}$. In Section 2, we review the notion of accessible singularity and local index. In Section 3, we compare the systems (1) with other types of coupled Painlevé II systems in dimension 4. In Section 4, we will prove Theorem 0.1 by giving an explicit birational transformation for each step. In the final section, we will prove Theorem 0.2.

1. RELATION BETWEEN THE SYSTEMS (1) AND THE SYSTEMS OF TYPE $A_4^{(1)}$

As is well-known, the degeneration from P_{IV} to P_{II} is given by

$$\beta_1 = \frac{1}{4\epsilon^6}, \quad \beta_2 = \alpha_1, \quad t = \frac{-1 + \epsilon^4 T}{\sqrt{2}\epsilon^3}, \quad x = \frac{1 + 2\epsilon^2 X}{\sqrt{2}\epsilon^3}, \quad y = \frac{\epsilon Y}{\sqrt{2}}.$$

Here the change of variables from (x, y) to (X, Y) is symplectic. As the fourth order analogue of the above confluence process, we consider the following degeneration from the systems of type $A_4^{(1)}$ to the systems (1).

The systems of type $A_4^{(1)}$ are given as follows:

$$\begin{cases} \frac{dx}{dt} = -x^2 + 4xy - 2tx - 2\beta_1 + 4zw \\ \frac{dy}{dt} = -2y^2 + 2xy + 2ty + \beta_2 \\ \frac{dz}{dt} = -z^2 + 4zw - 2tz - 2\beta_3 + 4yz \\ \frac{dw}{dt} = -2w^2 + 2zw + 2tw + \beta_4 - 4yw. \end{cases}$$

The systems of type $A_4^{(1)}$ are reduced to the systems (1) by putting

$$\begin{aligned} \beta_1 &= \frac{1}{4\epsilon^6}, \quad \beta_2 = \alpha_1, \quad \beta_3 = \frac{1}{4\epsilon^6}, \quad \beta_4 = \alpha_2, \quad t = \frac{-1 + \epsilon^4 T}{\sqrt{2}\epsilon^3}, \\ x &= \frac{1 + 2\epsilon^2 X}{\sqrt{2}\epsilon^3}, \quad y = \frac{\epsilon Y}{\sqrt{2}}, \quad z = \frac{1 + 2\epsilon^2 Z}{\sqrt{2}\epsilon^3}, \quad w = \frac{\epsilon W}{\sqrt{2}}, \end{aligned}$$

and taking the limit $\epsilon \rightarrow 0$.

2. ACCESSIBLE SINGULARITY

Let us review the notion of accessible singularity. Let B be a connected open domain in $\mathbb{C}$ and $\pi : \mathcal{W} \rightarrow B$ a smooth proper holomorphic map. We assume that $\mathcal{H} \subset \mathcal{W}$ is a normal crossing divisor which is flat over B . Let us consider a rational vector field $\tilde{v}$ on $\mathcal{W}$ satisfying the condition

$$\tilde{v} \in H^0(\mathcal{W}, \Theta_{\mathcal{W}}(-\log \mathcal{H})(\mathcal{H})).$$

Fixing $t_0 \in B$ and $P \in \mathcal{W}_{t_0}$, we can take a local coordinate system $(x_1, x_2, \dots, x_n)$ of $\mathcal{W}_{t_0}$ centered at P such that $\mathcal{H}_{smooth}$ can be defined by the local equation $x_1 = 0$. Since $\tilde{v} \in H^0(\mathcal{W}, \Theta_{\mathcal{W}}(-\log \mathcal{H})(\mathcal{H}))$, we can write down the vector field $\tilde{v}$ near $P = (0, 0, \dots, 0, t_0)$ as follows:

$$\tilde{v} = \frac{\partial}{\partial t} + a_1 \frac{\partial}{\partial x_1} + \frac{a_2}{x_1} \frac{\partial}{\partial x_2} + \dots + \frac{a_n}{x_1} \frac{\partial}{\partial x_n}$$

This vector field defines the following system of differential equations

$$(2) \quad \begin{cases} \frac{dx_1}{dt} = a_1(x_1, x_2, \dots, x_n, t) \\ \frac{dx_2}{dt} = \frac{a_2(x_1, x_2, \dots, x_n, t)}{x_1} \\ \cdot \\ \cdot \\ \cdot \\ \frac{dx_n}{dt} = \frac{a_n(x_1, x_2, \dots, x_n, t)}{x_1}. \end{cases}$$

Here $a_i(x_1, x_2, \dots, x_n, t)$ ($i = 1, 2, \dots, n$) are holomorphic functions near $P = (0, \dots, 0, t_0)$,

Definition 2.1. *With the above notation, assume that the rational vector field $\tilde{v}$ on $\mathcal{W}$ satisfies the condition*

$$\tilde{v} \in H^0(\mathcal{W}, \Theta_{\mathcal{W}}(-\log \mathcal{H})(\mathcal{H})).$$

We say that $\tilde{v}$ has an accessible singularity at $P = (0, 0, \dots, 0, t_0)$ if

$$x_1 = 0 \text{ and } a_i(0, 0, \dots, 0, t_0) = 0 \text{ for every } i, \ 2 \leq i \leq n.$$

If $P \in \mathcal{H}_{smooth}$ is not an accessible singularity, all solutions of the ordinary differential equation passing through P are vertical solutions, that is, the solutions are contained in the fiber $\mathcal{W}_{t_0}$ over $t = t_0$. If $P \in \mathcal{H}_{smooth}$ is an accessible singularity, there may be a solution of (2) which passes through P and goes into the interior $\mathcal{W} - \mathcal{H}$ of $\mathcal{W}$.

Let us recall the notion of local index. When we construct the phase spaces of the higher order Painlevé equations, an object that we call the local index is the key for determining when we need to make a blowing-up of an accessible singularity or a blowing-down to a minimal phase space. In the case of equations of higher order with favorable properties, for example the systems of type $A_4^{(1)}$ [1], the local index at the accessible singular point corresponds to the set of orders that appears in the free parameters of formal solutions passing through that point [9].

Definition 2.2. *Let v be an algebraic vector field which is given by (2) and (X, Y, Z, W) be a boundary coordinate system in a neighborhood of an accessible singularity $P = (0, 0, 0, 0, t)$. Assume that the system is written as*

$$\left\{ \begin{array}{l} \frac{dX}{dt} = a + f_1(X, Y, Z, W, t) \\ \frac{dY}{dt} = \frac{bY + f_2(X, Y, Z, W, t)}{X} \\ \frac{dZ}{dt} = \frac{cZ + f_3(X, Y, Z, W, t)}{X} \\ \frac{dW}{dt} = \frac{dW + f_4(X, Y, Z, W, t)}{X} \end{array} \right.$$

near the accessible singularity P , where a, b, c and d are nonzero constants. We say that the vector field v has the local index (a, b, c, d) at P if $f_1(X, Y, Z, W, t)$ is a polynomial which vanishes at $P = (0, 0, 0, 0, t)$ and $f_i(X, Y, Z, W, t)$ ($i = 2, 3, 4$) are polynomials of order 2 in X, Y, Z, W . Here $f_i \in \mathbb{C}[X, Y, Z, W, t]$ ($i = 1, 2, 3, 4$).

Remark 2.1. We are interested in the case with local index $(1, \frac{b}{a}, \frac{c}{a}, \frac{d}{a}) \in \mathbb{Z}^4$. If each component of $(1, \frac{b}{a}, \frac{c}{a}, \frac{d}{a})$ has the same sign, we may resolve the accessible singularity by blowing-up finitely many times. But when different signs appear, we may need to both blow up and blow down.

3. COMPARE THE SYSTEMS (1) WITH OTHER TYPES OF COUPLED PAINLEVÉ II SYSTEMS IN DIMENSION 4

It is known that certain reduction of the Drinfeld-Sokolov hierarchy of type $C_2^{(1)}$ reduces to a fourth order differential system with affine Weyl group symmetry

$W(C_2^{(1)})$ [2]. This system is an autonomous ordinary differential equations for 5 unknown functions f_0, f_1, f_2, u_1, u_2 involving complex parameters $\alpha_0, \alpha_1, \alpha_2$ satisfying $\alpha_0 + 2\alpha_1 + \alpha_2 = -4$, which are given as follows:

$$\begin{cases} \frac{df_0}{dT} = -2u_1f_0 - \alpha_0 \\ \frac{df_1}{dT} = (u_1 - u_2)f_1 - \alpha_1 \\ \frac{df_2}{dT} = 2u_2f_2 - \alpha_2 \\ \frac{du_1}{dT} = (u_1 + u_2)u_1 - \frac{f_1 - f_2}{h} \\ \frac{du_2}{dT} = -(u_1 + u_2)u_2 - \frac{f_0 - f_1}{h}, \end{cases}$$

where $h' = 0$, $f_0 + 2f_1 + f_2 = 4T + 2hu_1u_2$. Setting the variables x, y, z, w, t and the nonzero parameter h as

$$x = \frac{-u_1}{2}, \quad y = \frac{f_0}{8}, \quad z = \frac{u_2}{2}, \quad w = \frac{f_2}{8}, \quad t = 2T, \quad h = -1,$$

we can obtain the following coupled Painlevé II systems in dimension 4:

$$(3) \quad \begin{cases} \frac{dx}{dt} = -x^2 + y + 3w - \frac{t}{4} \\ \frac{dy}{dt} = 2xy + \alpha_1 \\ \frac{dz}{dt} = -z^2 + 3y + w - \frac{t}{4} \\ \frac{dw}{dt} = 2zw + \alpha_2. \end{cases}$$

Here the Hamiltonian of this system is given as follows:

$$\begin{aligned} H &= \frac{y^2}{2} - (x^2 + \frac{t}{4})y - \alpha_1x + \frac{w^2}{2} - (z^2 + \frac{t}{4})w - \alpha_2z + 3yw \\ &= H_{II}(x, y, t; \alpha_1) + H_{II}(z, w, t; \alpha_2) + 3yw. \end{aligned}$$

From the viewpoint of the local index, there are the following differences between these coupled Painlevé II systems.

Notation

$$\cdot (X, Y, Z, W) = (\frac{y}{x^2} - 2, \frac{1}{x}, \frac{z}{x}, \frac{w}{x}).$$

Systems	Accessible singularities	Type of local index
(1)	(X,Y,Z,W)=(0,0,1,0)	(-4,-1,-3,+1)
	(X,Y,Z,W)=(0,0,-2,0)	(-4,-1,+3,-5)
(3)	(X,Y,Z,W)=(0,0,2,0)	(-4,-1,-5,+3)
	(X,Y,Z,W)=(0,0,-3,0)	(-4,-1,+5,-7)
$P_{II} \times P_{II}$	(X,Y,Z,W)=(0,0,0,0)	(-4,-1,-1,-1)
	(X,Y,Z,W)=(0,0,-1,0)	(-4,-1,+1,-3)

Remark 3.1. *I don't know whether the accessible singular points $(X, Y, Z, W) = (0, 0, 2, 0)$ and $(X, Y, Z, W) = (0, 0, -3, 0)$ of the systems (3) can be resolved or not.*

4. PROOF OF THEOREM 0.1.

In this section, we will give an explicit resolution of accessible singularities of the systems (1) and will construct a family of phase spaces for the systems. In the case of Painlevé equations, we can obtain their phase spaces by adding subspaces of codimension 1 to the original space. But in the case of fourth order differential equations, we need to add codimension 2 subvarieties to the original space in addition to codimension 1 subvarieties.

4.1. Accessible singularities of the systems (1)

Let P be an accessible singular point in the boundary divisor $\mathcal{H}$ and (X, Y, Z, W) a coordinate system centered at P , where $\{X = 0\} \subset \mathcal{H}$. Rewriting the systems in the local coordinate system (X, Y, Z, W) , the right hand side of each differential equation has poles along $\mathcal{H}$. If we resolve the accessible singularity P and the right hand side of each differential equation becomes holomorphic in the coordinate system $(X', Y', Z', W') \in U \cong \mathbb{C}^4$, then we can use Cauchy's existence and uniqueness theorem of solutions. In order to consider a family of phase spaces for the system (1), let us take the compactification $\mathbb{P}^4 \times B$ of $\mathbb{C}^4 \times B$. Moreover, we denote the boundary divisor in $\mathbb{P}^4$ by $\mathcal{H}$. Fixing the parameter α_i , consider the product $\mathbb{P}^4 \times B$ and extend the regular vector field on $\mathbb{C}^4 \times B$ to a rational vector field $\tilde{v}$ on $\mathbb{P}^4 \times B$. The following Lemma 4.1 shows that this rational vector field $\tilde{v}$ has six accessible singular points on the boundary divisor $\mathcal{H} \times \{t\} \subset \mathbb{P}^4 \times \{t\}$ for each $t \in B$.




FIGURE 1

Lemma 4.1. *The rational vector field $\tilde{v}$ has six accessible singular points $P_i = \{(X_i, Y_i, Z_i, W_i) = (0, 0, 0, 0)\}$ ($i = 1, 2, 3, 4$) and $P_5 = \{(X_1, Y_1, Z_1, W_1) | X_1 = Y_1 = W_1 = 0, Z_1 = 1\}$ and $P_6 = \{(X_3, Y_3, Z_3, W_3) | X_3 = Y_3 = Z_3 = 0, W_3 = -1\}$. Here, $(X_1, Y_1, Z_1, W_1) = (\frac{1}{x}, \frac{y}{x}, \frac{z}{x}, \frac{w}{x})$, $(X_2, Y_2, Z_2, W_2) = (\frac{x}{z}, \frac{y}{z}, \frac{1}{z}, \frac{w}{z})$, $(X_3, Y_3, Z_3, W_3) = (\frac{x}{y}, \frac{1}{y}, \frac{z}{y}, \frac{w}{y})$ and $(X_4, Y_4, Z_4, W_4) = (\frac{x}{w}, \frac{y}{w}, \frac{z}{w}, \frac{1}{w})$.*

This lemma can be proven by direct calculation. □

Remark 4.1. *By the symmetry*

$$\pi : (x, y, z, w; \alpha_1, \alpha_2, \alpha_3) \longrightarrow (z, w, x, y; \alpha_1, \alpha_3, \alpha_2),$$

it is easy to see that

$$\pi(P_1) = P_2, \quad \pi(P_3) = P_4.$$

Now we are ready to prove Theorem 0.1.

4.2. Resolution of the accessible singular point P_1

In this subsection, we will give an explicit resolution process for the accessible singular point P_1 by giving a convenient coordinate system at each step.

By the following steps, we can resolve the accessible singular point P_1 .

Step 1: We blow up at the point P_1

$$x_1^{(1)} = X_1, \quad y_1^{(1)} = \frac{Y_1}{X_1}, \quad z_1^{(1)} = \frac{Z_1}{X_1}, \quad w_1^{(1)} = \frac{W_1}{X_1}.$$

Step 2: We blow up along the surface $\{(x_1^{(1)}, y_1^{(1)}, z_1^{(1)}, w_1^{(1)}) | x_1^{(1)} = y_1^{(1)} = 0\}$

$$x_1^{(2)} = x_1^{(1)}, \quad y_1^{(2)} = \frac{y_1^{(1)}}{x_1^{(1)}}, \quad z_1^{(2)} = z_1^{(1)}, \quad w_1^{(2)} = w_1^{(1)}.$$

Step 3: We blow up along the surface $\{(x_1^{(2)}, y_1^{(2)}, z_1^{(2)}, w_1^{(2)}) | y_1^{(2)} - \alpha_1 = x_1^{(2)} = 0\}$

$$x_1^{(3)} = x_1^{(2)}, \quad y_1^{(3)} = \frac{y_1^{(2)} - \alpha_1}{x_1^{(2)}}, \quad z_1^{(3)} = z_1^{(2)}, \quad w_1^{(3)} = w_1^{(2)}.$$

4.3. Resolution of the accessible singular point P_5

In this subsection, we will give an explicit resolution process for the accessible singular point P_5 by giving a convenient coordinate system at each step.

By the following steps, we can resolve the accessible singular point P_5 . At first, we take the coordinate system $\{(X_5, Y_5, Z_5, W_5) = (X_1, Y_1, Z_1 - 1, W_1)\}$ centered at P_5 .

Step 1: We blow up along the curve $\{(X_5, Y_5, Z_5, W_5) | X_5 = Y_5 = W_5 = 0\} \cong \mathbb{P}^1$

$$x_5^{(1)} = X_5, \quad y_5^{(1)} = \frac{Y_5}{X_5}, \quad z_5^{(1)} = Z_5, \quad w_5^{(1)} = \frac{W_5}{X_5}.$$

Step 2: We blow down the 3-fold $\{(x_5^{(1)}, y_5^{(1)}, z_5^{(1)}, w_5^{(1)}) | x_5^{(1)} = 0\} \cong \mathbb{P}^2 \times \mathbb{P}^1$

$$x_5^{(2)} = x_5^{(1)}, \quad y_5^{(2)} = y_5^{(1)}, \quad z_5^{(2)} = \frac{x_5^{(1)}}{z_5^{(1)} + 1}, \quad w_5^{(2)} = w_5^{(1)}.$$




FIGURE 2. $\mathbb{P}^2$ -flop

The resolution process from Step 1 to Step 2 is well-known as $\mathbb{P}^2$ -flop. In order to resolve the accessible singular point P_5 and obtain a holomorphic coordinate system, we need to blow down the 3-fold $V_1 \cong \mathbb{P}^2 \times \mathbb{P}^1$ along the $\mathbb{P}^1$ -fiber. After we blow down the 3-fold V_1 , we can resolve the accessible singular point P_5 by only blowing-ups.

Step 3: We blow up along the surfaces $\{(x_5^{(2)}, y_5^{(2)}, z_5^{(2)}, w_5^{(2)}) | x_5^{(2)} = y_5^{(2)} = 0\}$ and $\{(x_5^{(2)}, y_5^{(2)}, z_5^{(2)}, w_5^{(2)}) | w_5^{(2)} = z_5^{(2)} = 0\}$

$$x_5^{(3)} = x_5^{(2)}, \quad y_5^{(3)} = \frac{y_5^{(2)}}{x_5^{(2)}}, \quad z_5^{(3)} = z_5^{(2)}, \quad w_5^{(3)} = \frac{w_5^{(2)}}{z_5^{(2)}}.$$

Step 4: We blow up along the surfaces $\{(x_5^{(3)}, y_5^{(3)}, z_5^{(3)}, w_5^{(3)}) | x_5^{(3)} = y_5^{(3)} - \alpha_1 = 0\}$ and $\{(x_5^{(3)}, y_5^{(3)}, z_5^{(3)}, w_5^{(3)}) | w_5^{(3)} - \alpha_2 = z_5^{(3)} = 0\}$

$$x_5^{(4)} = x_5^{(3)}, \quad y_5^{(4)} = \frac{y_5^{(3)} - \alpha_1}{x_5^{(3)}}, \quad z_5^{(4)} = z_5^{(3)}, \quad w_5^{(4)} = \frac{w_5^{(3)} - \alpha_2}{z_5^{(3)}}.$$

4.4. Resolution of the accessible singular point P_6

In this subsection, we will give an explicit resolution process for the accessible singular point P_6 by giving a convenient coordinate system at each step.

By the following steps, we can resolve the accessible singular point P_6 . At first, we take the coordinate system $(x_8^{(0)}, y_8^{(0)}, z_8^{(0)}, w_8^{(0)})$ centered at P_6 .

Step 1: We blow up at the point P_6

$$x_8^{(1)} = \frac{x_8^{(0)}}{y_8^{(0)}}, \quad y_8^{(1)} = y_8^{(0)}, \quad z_8^{(1)} = \frac{z_8^{(0)}}{y_8^{(0)}}, \quad w_8^{(1)} = \frac{w_8^{(0)}}{y_8^{(0)}}.$$

Step 2: We blow up along the surface $\{(x_8^{(1)}, y_8^{(1)}, z_8^{(1)}, w_8^{(1)}) | x_8^{(1)} - z_8^{(1)} = y_8^{(1)} = 0\}$

$$x_8^{(2)} = \frac{x_8^{(1)} - z_8^{(1)}}{y_8^{(1)}}, \quad y_8^{(2)} = y_8^{(1)}, \quad z_8^{(2)} = z_8^{(1)}, \quad w_8^{(2)} = w_8^{(1)}.$$

Step 3: We blow up along the surface $\{(x_8^{(2)}, y_8^{(2)}, z_8^{(2)}, w_8^{(2)}) | x_8^{(2)} + \alpha_1 + \alpha_2 - 1 = y_8^{(2)} = 0\}$

$$x_8^{(3)} = \frac{x_8^{(2)} + \alpha_1 + \alpha_2 - 1}{y_8^{(2)}}, \quad y_8^{(3)} = y_8^{(2)}, \quad z_8^{(3)} = z_8^{(2)}, \quad w_8^{(3)} = w_8^{(2)}.$$

4.5. Resolution of the accessible singular point P_3

In this subsection, we will give an explicit resolution process for the accessible singular point P_3 by giving a convenient coordinate system at each step.

By the following steps, we can resolve the accessible singular point P_3 .

Step 1: We blow up along the curve $\{(X_3, Y_3, Z_3, W_3) | X_3 = Y_3 = Z_3 = 0\}$

$$x_3^{(1)} = X_3, \quad y_3^{(1)} = \frac{Y_3}{X_3}, \quad z_3^{(1)} = \frac{Z_3}{X_3}, \quad w_3^{(1)} = W_3.$$

Step 2: We blow up along the surface $\{(x_3^{(1)}, y_3^{(1)}, z_3^{(1)}, w_3^{(1)}) | y_3^{(1)} = x_3^{(1)} = 0\}$

$$x_3^{(2)} = \frac{x_3^{(1)}}{y_3^{(1)}} , \quad y_3^{(2)} = y_3 , \quad z_3^{(2)} = z_3^{(1)} , \quad w_3^{(2)} = w_3^{(1)} .$$

Step 3: We blow up along the surface $\{(x_3^{(2)}, y_3^{(2)}, z_3^{(2)}, w_3^{(2)}) | x_3^{(2)} = w_3^{(2)} = 0\}$

$$x_3^{(3)} = x_3^{(2)} , \quad y_3^{(3)} = y_3^{(2)} , \quad z_3^{(3)} = z_3^{(2)} , \quad w_3^{(3)} = \frac{w_3^{(2)}}{x_3^{(2)}} .$$

Step 4: We make a change of variables

$$x_3^{(4)} = \frac{1}{x_3^{(3)}} , \quad y_3^{(4)} = y_3^{(3)} , \quad z_3^{(4)} = z_3^{(3)} , \quad w_3^{(4)} = w_3^{(3)} .$$

This change of variables is necessary for making the transition functions in the description of $\mathcal{X}$ symplectic [3]. It is easy to see that there are two accessible singular points $P_3 = \{(x_3^{(4)}, y_3^{(4)}, z_3^{(4)}, w_3^{(4)}) | x_3^{(4)} + w_3^{(4)} = 2, y_3^{(4)} = w_3^{(4)} = 0, z_3^{(4)} = 1\}$ and $P_6 = \{(x_3^{(4)}, y_3^{(4)}, z_3^{(4)}, w_3^{(4)}) | x_3^{(4)} + w_3^{(4)} = 2, y_3^{(4)} = w_3^{(4)} = 0, z_3^{(4)} = -2\}$ in the domain $\{(x_3^{(4)}, y_3^{(4)}, z_3^{(4)}, w_3^{(4)}) | y_3^{(4)} = 0\} \cong \mathbb{C}^3$.

Step 5: We blow up along the surface $\{(x_3^{(4)}, y_3^{(4)}, z_3^{(4)}, w_3^{(4)}) | y_3^{(4)} = z_3^{(4)} - 1 = 0\}$

$$x_3^{(5)} = x_3^{(4)} , \quad y_3^{(5)} = y_3^{(4)} , \quad z_3^{(5)} = \frac{z_3^{(4)} - 1}{y_3^{(4)}} , \quad w_3^{(5)} = w_3^{(4)} .$$

Step 6: We blow up along the surface $\{(x_3^{(5)}, y_3^{(5)}, z_3^{(5)}, w_3^{(5)}) | x_3^{(5)} + w_3^{(5)} - 2 = y_3^{(5)} = 0\}$

$$x_3^{(6)} = \frac{x_3^{(5)} + w_3^{(5)} - 2}{y_3^{(5)}} , \quad y_3^{(6)} = y_3^{(5)} , \quad z_3^{(6)} = z_3^{(5)} , \quad w_3^{(6)} = w_3^{(5)} .$$

Step 7: We blow up along the surface $\{(x_3^{(6)}, y_3^{(6)}, z_3^{(6)}, w_3^{(6)}) | y_3^{(6)} = z_3^{(6)} = 0\}$

$$x_3^{(7)} = x_3^{(6)} , \quad y_3^{(7)} = y_3^{(6)} , \quad z_3^{(7)} = \frac{z_3^{(6)}}{y_3^{(6)}} , \quad w_3^{(7)} = w_3^{(6)} .$$

Step 8: We blow up along the surface $\{(x_3^{(7)}, y_3^{(7)}, z_3^{(7)}, w_3^{(7)}) | x_3^{(7)} = y_3^{(7)} = 0\}$

$$x_3^{(8)} = \frac{x_3^{(7)}}{y_3^{(7)}} , \quad y_3^{(8)} = y_3^{(7)} , \quad z_3^{(8)} = z_3^{(7)} , \quad w_3^{(8)} = w_3^{(7)} .$$

Step 9: We blow up along the surface $\{(x_3^{(8)}, y_3^{(8)}, z_3^{(8)}, w_3^{(8)}) | y_3^{(8)} = z_3^{(8)} = 0\}$

$$x_3^{(9)} = x_3^{(8)} , \quad y_3^{(9)} = y_3^{(8)} , \quad z_3^{(9)} = \frac{z_3^{(8)}}{y_3^{(8)}} , \quad w_3^{(9)} = w_3^{(8)} .$$

Step 10: We blow up along the surface $\{(x_3^{(9)}, y_3^{(9)}, z_3^{(9)}, w_3^{(9)}) | x_3^{(9)} - t = y_3^{(9)} = 0\}$

$$x_3^{(10)} = \frac{x_3^{(9)} - t}{y_3^{(9)}} , \quad y_3^{(10)} = y_3^{(9)} , \quad z_3^{(10)} = z_3^{(9)} , \quad w_3^{(10)} = w_3^{(9)} .$$

Step 11: We blow up along the surface $\{(x_3^{(10)}, y_3^{(10)}, z_3^{(10)}, w_3^{(10)}) | y_3^{(10)} = x_3^{(10)} - 2z_3^{(10)}w_3^{(10)} + 1 - \alpha_1 - \alpha_2 = 0\}$

$$x_3^{(11)} = \frac{x_3^{(10)} - 2z_3^{(10)}w_3^{(10)} + 1 - \alpha_1 - \alpha_2}{y_3^{(10)}}, \quad y_3^{(11)} = y_3^{(10)}, \quad z_3^{(11)} = z_3^{(10)}, \quad w_3^{(11)} = w_3^{(10)}.$$

4.6. Resolution of the accessible singular point P_6

In this subsection, we will give an explicit resolution process for the accessible singular point P_6 by giving a convenient coordinate system at each step.

By the following steps, we can resolve the accessible singular point P_6 .

Step 1: We take the coordinate system centered at P_6

$$x_6^{(1)} = x_3^{(4)} + w_3^{(4)} - 2, \quad y_6^{(1)} = y_3^{(4)}, \quad z_6^{(1)} = z_3^{(4)} + 2, \quad w_6^{(1)} = w_3^{(4)}.$$

Step 2: We blow down the 3-fold $\{(x_6^{(1)}, y_6^{(1)}, z_6^{(1)}, w_6^{(1)}) | y_6^{(1)} = 0\} \subset \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$

$$x_6^{(2)} = x_6^{(1)}, \quad y_6^{(2)} = y_6^{(1)}, \quad z_6^{(2)} = \frac{y_6^{(1)}z_6^{(1)}}{1 - \frac{z_6^{(1)}}{3}}, \quad w_6^{(2)} = w_6^{(1)}.$$

Step 3: We make a change of variables

$$x_6^{(3)} = x_6^{(2)}, \quad y_6^{(3)} = y_6^{(2)}, \quad z_6^{(3)} = z_6^{(2)} + 3y_6^{(2)}, \quad w_6^{(3)} = w_6^{(2)}.$$

Step 4: We blow up along the surface $\{(x_6^{(3)}, y_6^{(3)}, z_6^{(3)}, w_6^{(3)}) | x_6^{(3)} = y_6^{(3)} = 0\}$

$$x_6^{(4)} = \frac{x_6^{(3)}}{y_6^{(3)}}, \quad y_6^{(4)} = y_6^{(3)}, \quad z_6^{(4)} = z_6^{(3)}, \quad w_6^{(4)} = w_6^{(3)}.$$

Step 5: We blow down the 3-fold $\{(x_6^{(4)}, y_6^{(4)}, z_6^{(4)}, w_6^{(4)}) | y_6^{(4)} = 0\} \subset \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$

$$x_6^{(5)} = x_6^{(4)}, \quad y_6^{(5)} = y_6^{(4)}, \quad z_6^{(5)} = y_6^{(4)}z_6^{(4)}, \quad w_6^{(5)} = w_6^{(4)}.$$

Step 6: We blow up along the surface $\{(x_6^{(5)}, y_6^{(5)}, z_6^{(5)}, w_6^{(5)}) | x_6^{(5)} = y_6^{(5)} = 0\}$

$$x_6^{(6)} = \frac{x_6^{(5)}}{y_6^{(5)}}, \quad y_6^{(6)} = y_6^{(5)}, \quad z_6^{(6)} = z_6^{(5)}, \quad w_6^{(6)} = w_6^{(5)}.$$

Step 7: We blow down the 3-fold $\{(x_6^{(6)}, y_6^{(6)}, z_6^{(6)}, w_6^{(6)}) | y_6^{(6)} = 0\} \subset \mathbb{P}^1 \times \mathbb{P}^1 \times \mathbb{P}^1$

$$x_6^{(7)} = x_6^{(6)}, \quad y_6^{(7)} = y_6^{(6)}, \quad z_6^{(7)} = y_6^{(6)}z_6^{(6)}, \quad w_6^{(7)} = w_6^{(6)}.$$

Step 8: We blow up along the surfaces $\{(x_6^{(7)}, y_6^{(7)}, z_6^{(7)}, w_6^{(7)}) | x_6^{(7)} - t = y_6^{(7)} = 0\}$ and $\{(x_6^{(7)}, y_6^{(7)}, z_6^{(7)}, w_6^{(7)}) | z_6^{(7)} = w_6^{(7)} = 0\}$

$$x_6^{(8)} = \frac{x_6^{(7)} - t}{y_6^{(7)}}, \quad y_6^{(8)} = y_6^{(7)}, \quad z_6^{(8)} = z_6^{(7)}, \quad w_6^{(8)} = \frac{w_6^{(7)}}{z_6^{(7)}}.$$

Step 9: We blow up along the surfaces $\{(x_6^{(8)}, y_6^{(8)}, z_6^{(8)}, w_6^{(8)}) | x_6^{(8)} + 18w_6^{(8)} + 1 - \alpha_1 - \alpha_2 = y_6^{(8)} = 0\}$ and $\{(x_6^{(8)}, y_6^{(8)}, z_6^{(8)}, w_6^{(8)}) | w_6^{(8)} - \frac{\alpha_2}{9} = z_6^{(8)} = 0\}$

$$x_6^{(9)} = \frac{x_6^{(8)} + 18w_6^{(8)} + 1 - \alpha_1 - \alpha_2}{y_6^{(8)}}, \quad y_6^{(9)} = y_6^{(8)}, \quad z_6^{(9)} = z_6^{(8)}, \quad w_6^{(9)} = \frac{w_6^{(8)} - \frac{\alpha_2}{9}}{z_6^{(8)}}.$$

5. HAMILTONIAN SYSTEM

The system (1) is written as a Hamiltonian system:

$$\frac{dx}{dt} = \frac{\partial H}{\partial y}, \quad \frac{dy}{dt} = -\frac{\partial H}{\partial x}, \quad \frac{dz}{dt} = \frac{\partial H}{\partial w}, \quad \frac{dw}{dt} = -\frac{\partial H}{\partial z}.$$

Here the Hamiltonian is given as follows:

$$H = \frac{y^2}{2} - (x^2 + \frac{t}{2})y - \alpha_1 x + \frac{w^2}{2} - (z^2 + \frac{t}{2})w - \alpha_2 z + yw.$$

We list below the Hamiltonian for each coordinate system (x_i, y_i, z_i, w_i) ($i = 1, 3, 5, 6$).

The Hamiltonian H_1 in U_1 We obtain

$$H_1 = \frac{1}{2}(-tw_1 + w_1^2 + 2y_1 + tx_1^2y_1 - 2x_1^2y_1w_1 + x_1^4y_1^2 - 2z_1^2w_1 + t\alpha_1x_1 - 2\alpha_1x_1w_1 + 2\alpha_1x_1^3y_1 + \alpha_1^2x_1^2 - 2\alpha_2z_1),$$

where

$$x_1 = \frac{1}{x}, \quad y_1 = -x(xy + \alpha_1), \quad z_1 = z, \quad w_1 = w.$$

The Hamiltonian H_1 and coordinate system (x_1, y_1, z_1, w_1) above satisfy the following condition:

$$dx_1 \wedge dy_1 + dz_1 \wedge dw_1 - dH_1 \wedge dt = dx \wedge dy + dz \wedge dw - dH \wedge dt.$$

The Hamiltonian H_3 in U_3 We obtain

$$H_3 = \frac{1}{2}(2x_3 - ty_3 + y_3^2 + tx_3y_3^2 - 2x_3y_3^3 + x_3^2y_3^4 + 2ty_3z_3w_3 - 4y_3^2z_3w_3 + 4x_3y_3^3z_3w_3 - 2y_3^2z_3^2w_3 + 4y_3^2z_3^2w_3^2 + t\alpha_1y_3 - 2\alpha_1y_3^2 + 2\alpha_1x_3y_3^3 + 4\alpha_1y_3^2z_3w_3 + \alpha_1^2y_3^2 + t\alpha_2y_3 - 2\alpha_2y_3^2 + 2\alpha_2x_3y_3^3 - 2\alpha_2y_3^2z_3 + 4\alpha_2y_3^2z_3w_3 + 2\alpha_1\alpha_2y_3^2 + \alpha_2^2y_3^2),$$

where

$$x_3 = x(1 + (y + w - 2x^2 - t)x - 2w(z - x) - \alpha_1 - \alpha_2), \quad y_3 = \frac{1}{x}, \quad z_3 = x^2(z - x), \quad w_3 = \frac{w}{x^2}.$$

The Hamiltonian H_3 and coordinate system (x_3, y_3, z_3, w_3) above satisfy the following condition:

$$dx_3 \wedge dy_3 + dz_3 \wedge dw_3 - dH_3 \wedge dt = dx \wedge dy + dz \wedge dw - d(H + x) \wedge dt.$$

The Hamiltonian H_5 in U_5 We obtain

$$H_5 = \frac{1}{2}(2w_5 + 2y_5 + tx_5^2y_5 + x_5^4y_5^2 + tz_5^2w_5 + 2x_5^2y_5z_5^2w_5 + z_5^4w_5^2 + t\alpha_1x_5 + 2\alpha_1x_5^3y_5 \\ + 2\alpha_1x_5z_5^2w_5 + \alpha_1^2x_5^2 + t\alpha_2z_5 + 2\alpha_2x_5^2y_5z_5 + 2\alpha_2z_5^3w_5 + 2\alpha_1\alpha_2x_5z_5 + \alpha_2^2z_5^2),$$

where

$$x_5 = \frac{1}{x}, \quad y_5 = -x(xy + \alpha_1), \quad z_5 = \frac{1}{z}, \quad w_5 = -z(zw + \alpha_2).$$

The Hamiltonian H_5 and coordinate system (x_5, y_5, z_5, w_5) above satisfy the following condition:

$$dx_5 \wedge dy_5 + dz_5 \wedge dw_5 - dH_5 \wedge dt = dx \wedge dy + dz \wedge dw - dH \wedge dt.$$

The Hamiltonian H_6 in U_6 We obtain

$$H_6 = \frac{1}{2}(2x_6 - ty_6 + y_6^2 - 162y_6^2w_6 + tx_6y_6^2 - 2x_6y_6^3 + x_6^2y_6^4 - 2ty_6z_6w_6 \\ + 4y_6^2z_6w_6 - 4x_6y_6^3z_6w_6 + 4y_6^2z_6^2w_6^2 + t\alpha_1y_6 - 2\alpha_1y_6^2 + 2\alpha_1x_6y_6^3 - 4\alpha_1y_6^2z_6w_6 \\ + \alpha_1^2y_6^2 - t\alpha_2y_6 + 2\alpha_2y_6^2 - 2\alpha_2x_6y_6^3 + 4\alpha_2y_6^2z_6w_6 - 2\alpha_1\alpha_2y_6^2 + \alpha_2^2y_6^2),$$

where

$$x_6 = x(1 + (y + w - 2x^2 - t)x + 2(xw - zw - \alpha_2) - \alpha_1 + \alpha_2), \quad y_6 = \frac{1}{x}, \\ z_6 = \frac{81}{x^2(x - z)}, \quad w_6 = \frac{x^2(x - z)(xw - zw - \alpha_2)}{81}.$$

The Hamiltonian H_6 and coordinate system (x_6, y_6, z_6, w_6) above satisfy the following condition:

$$dx_6 \wedge dy_6 + dz_6 \wedge dw_6 - dH_6 \wedge dt = dx \wedge dy + dz \wedge dw - d(H + x) \wedge dt.$$

The Hamiltonian H_8 in U_8 We obtain

$$H_8 = \frac{1}{2}(-tw_8 + w_8^2 - 2y_8 + 4x_8y_8^2 - 2x_8^2y_8^3 - 4z_8 + 4x_8y_8z_8 - 2z_8^2w_8 + 2\alpha_1y_8 \\ - 2\alpha_1x_8y_8^2 + 2\alpha_1z_8 + 4\alpha_2y_8 - 4\alpha_2x_8y_8^2 + 2\alpha_2z_8 - 2\alpha_1\alpha_2y_8 - 2\alpha_2^2y_8),$$

where

$$x_8 = -((x - z)y - 1 + \alpha_1 + \alpha_2)y, \quad y_8 = \frac{1}{y}, \quad z_8 = z, \quad w_8 = w + y.$$

The Hamiltonian H_8 and coordinate system (x_8, y_8, z_8, w_8) above satisfy the following condition:

$$dx_8 \wedge dy_8 + dz_8 \wedge dw_8 - dH_8 \wedge dt = dx \wedge dy + dz \wedge dw - dH \wedge dt.$$

Acknowledgement. The author thanks Professors K. Takano and Y. Yamada for giving helpful suggestions and encouragement, and Professor M. Noumi for giving his notes and helpful advice, and Professor W. Rossman for checking English.

REFERENCES

- [1] M. Noumi and Y. Yamada, *Higher order Painlevé equations of type $A_l^{(1)}$* , Funkcial. Ekvac., **41**, 1998, 483–503.
- [2] M. Noumi, *Private communication*.
- [3] T. Matano, A. Matsumiya and K. Takano, *On some Hamiltonian structures of Painlevé systems, II*, J. Math. Soc. Japan, **51**, No.4, 1999, 843–866.
- [4] T. Shioda and K. Takano, *On some Hamiltonian structures of Painlevé systems I*, Funkcial. Ekvac., **40**, 1997, 271–291.
- [5] Y. Sasano, *Coupled Painlevé IV systems in dimension 4*, preprint.
- [6] Y. Sasano, *Coupled Painlevé V systems in dimension 4*, preprint.
- [7] Y. Sasano, *Coupled Painlevé III systems in dimension 4 and the systems of type $A_5^{(1)}$* , preprint.
- [8] K. Takano, *Confluence processes in defining manifolds for Painlevé systems*, Tohoku Math. J. **53** (2001), 319–335.
- [9] N. Tahara, *An augmentation of the phase space of the system of type $A_4^{(1)}$* , Kyushu J. Math. **58**, 2004, 393–425.

Department of Mathematics
Kobe University
Kobe, Rokko, 657-8501
Japan

e-mail: sasano@math.kobe-u.ac.jp